

## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <a href="http://about.jstor.org/participate-jstor/individuals/early-journal-content">http://about.jstor.org/participate-jstor/individuals/early-journal-content</a>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

FOREIGN AND INSULAR STATISTICAL REPORTS OF COUNTRIES AND CITIES—UNTABULATED.

Australia—New South Wales—Newcastle.—Month of March, 1906. Estimated population, 51,050. Total number of deaths, 40, including 5 from tuberculosis.

France—St. Etienne.—Two weeks ended April 30, 1906. Estimated population, 146,836. Total number of deaths, 133, including enteric fever 1, scarlet fever 1, and 18 from tuberculosis.

GREAT BRITAIN—England and Wales.—The deaths registered in 76 great towns in England and Wales during the week ended May 12, 1906, correspond to an annual rate of 15.5 per 1,000 of population, which is estimated at 15,818,360.

London.—One thousand four hundred and seventeen deaths were registered during the week, including measles 75, scarlet fever 11, diphtheria 10, whooping cough 26, enteric fever 5, and 16 from diarrhea. The deaths from all causes correspond to an annual rate of 15.6 per 1,000. In Greater London 1,983 deaths were registered. In the "outer ring" the deaths included 4 from diphtheria, 9 from measles, 5 from scarlet fever, and 7 from whooping cough.

Ireland.—The average annual death rate represented by the deaths registered during the week ended May 12, 1906, in the 21 principal town districts of Ireland was 22.4 per 1,000 of the population, which is estimated at 1,093,959. The lowest rate was recorded in Ballymena, viz, 4.8, and the highest in Newtownards, viz, 45.8 per 1,000. In Dublin and suburbs 165 deaths were registered, including enteric fever 3, measles 1, whooping cough 1, and 28 from tuberculosis.

Scotland.—The deaths registered in 8 principal towns during the week ended May 12, 1906, correspond to an annual rate of 18.8 per 1,000 of the population, which is estimated at 1,787,788. The lowest rate of mortality was recorded in Paisley, viz, 13.2, and the highest in Perth, viz, 22.7 per 1,000. The aggregate number of deaths registered from all causes was 645, including diphtheria 7, enteric fever 1, measles 26, scarlet fever 2, and 16 from whooping cough.

Salford.—Three months ended March 31, 1906. Estimated population, 234,077. Total number of deaths, 976, including diphtheria 16, enteric fever 12, measles 8, scarlet fever 15, whooping cough 25, and 95 from phthisis pulmonalis.

Jamaica—Port Antonio.—Two weeks ended May 12, 1906. Estimated population, 3,000. Total number of deaths not reported. No contagious diseases reported.

Japan—Nagasaki.—Ten days ended April 30, 1906. Estimated population, 159,401. Total number of deaths not reported. No deaths from contagious diseases reported.

Malta.—Two weeks ended May 5, 1906. Estimated population,

633 June 8, 1906

Romarks

205,062. Total number of deaths, 153, including diphtheria 5 and 4 from whooping cough.

St. Helena.—Eight weeks ended April 28, 1906. Estimated population, 3,500. Total number of deaths, 7. No deaths from contagious diseases reported.

West Indies—Curaçoa.—Two weeks ended May 18, 1906. Estimated population, 31,600. Total number of deaths, 12. No contagious diseases.

Cholora, yellow fever, plague, and smallpox, from December 30, 1905, to June 8, 1906.

[Reports received by the Surgeon-General, Public Health and Marine-Hospital Service, from American consuls through the Department of State, and from other sources.]

[For reports received from June 30, 1905, to December 29, 1905, see Public Health Reports for December 29, 1905.]

[Note.—In accordance with custom, the tables of epidemic diseases are terminated semiannually and new tables begun.]

CHOLERA.

Date

Place

Cases. Deaths.

Place.	Date.	Cases.	Deaths.	Remarks.
India: Bombay Calcutta. Madras. Rangoon Philippine Islands: Provinces	Dec. 6-May 8 Nov. 12-Apr. 28 Nov. 25-Apr. 6 Dec. 17-Apr. 21	3,810	1,453 40 43 2,862	
Manila	Dec. 31-Apr. 21	283	252	
Russia: General	Nov. 23-Jan. 14 Dec. 4-29	34 72 29	16 11 34 12 5	
YELLOW FEVER.				
		·	1	
Brazil: Rio de Janeiro	Nov. 20-Apr. 22 Dec. 6-12		45 1	
Barranquilla Cartagena	Nov. 28-Dec. 4 Dec. 17-23		4	
Cuba: Habana Matanzas Province Ecuador:	Dec. 25-June 7 Feb. 5-May 23	}	6 2	
Guayaquil	Dec. 4-Apr. 30 Apr. 6-21		210	Estimate for March, 1906, about 300 cases, 60 deaths.
El Paraiso Puerto Cortez San Pedro	Jan. 13-19 Mar. 13 Mar. 18-24	<u>1</u>	1	Present.
Mexico: Oaxaca—				
Salina Cruz Tehuantepec Veracruz—	Mar. 18-24			Imported from Santa Lucrecia. Do.
Cordoba Orizaba Veracruz	Dec. 10–16 Dec. 10–Jan. 13 Dec. 24–Apr. 6	1 1 7	3 1 5	Imported from Quintana, Mex-
Yucatan— Merida	Dec. 17-May 19	20	8	iĉo.
Nicaragua: Managua	Dec. 10-Mar. 17		5	
Panama: Bocas del Toro Colon Peru:	Feb. 21 Dec. 8-May 22	2	1 1	
Callao	Mar. 4-Apr. 11	6	4	One on ss. Colombia from Guaya- quil, and 5 cases and 3 deaths on ss. Luxor.